

Mildly thick fluids (IDDSI level 2)

Mildly thick fluid looks like cream pouring from a bottle.

- It can be sipped.
- It can be drunk from a cup.
- It can only be drunk through a straw if a speech-language therapist recommends this.
- It falls quickly from a spoon but slower than water.

How to check a mildly thick fluid

A mildly thick fluid is the right consistency if it passes the IDDSI flow test. Ask a carer or friend to help you with this test.



IDDSI flow test

The IDDSI flow test measures the amount of fluid left in a syringe after 10 seconds of flow.

For a mildly thick fluid, 4 to 8 ml remains in the syringe.

See the IDDSI flow test videos at
www.iddsi.org/videos

Mildly thick fluids (IDDSI level 2)

Foods to avoid

- × Jelly.
- × Ice cream.
- × Canned or stewed fruit in juice.
- × Thin soup.
- × Thin sauces.
- × Porridge and cereals with fluid on top.

Notes or preparation tips

*Written by speech–language therapists, Te Whatu Ora – Health New Zealand Waitaha Canterbury. Adapted by HealthInfo clinical advisers.
Last reviewed July 2023.*

This is NOT an official IDDSI resource, educational material or education program and it is NOT meant to replace materials and resources on [IDDSI.org](https://iddsi.org). See [IDDSI.org](https://iddsi.org) for the most current information and resources. IDDSI descriptors © The International Dysphagia Diet Standardisation Initiative 2019 @ iddsi.org/framework/. Licensed under the Creative Commons Attribution Sharealike 4.0 License creativecommons.org/licenses/by-sa/4.0/legalcode. Derivative works extending beyond language translation are NOT PERMITTED.