

The National Immunisation Schedule

From – August 2025

Stages	Disease to protect against	Vaccine
Pregnancy	Tetanus + diphtheria + whooping cough (pertussis)	Boostrix®
	Influenza	Brand varies
6 weeks	Rotavirus	Rotarix® (oral) (first dose must be given before 15 weeks)
	Diphtheria + tetanus + whooping cough (pertussis) + polio + hepatitis B + Haemophilus influenzae type b (Hib)	Infanrix® hexa
	Pneumococcal disease	Prevenar 13®
3 months	Rotavirus	Rotarix® (oral) (second dose must be given before 25 weeks)
	Diphtheria + tetanus + whooping cough + polio + hepatitis B + Haemophilus influenzae type b (Hib)	Infanrix® hexa
	Meningococcal B	Bexsero® (can be given at 8 weeks)
5 months	Diphtheria + tetanus + whooping cough + polio + hepatitis B + Haemophilus influenzae type b (Hib)	Infanrix® hexa
	Pneumococcal disease	Prevenar 13®
	Meningococcal B	Bexsero® (can be given at 4 months)
12 months	Measles + mumps + rubella	Priorix®
	Pneumococcal disease	Prevenar 13®
	Meningococcal B	Bexsero®
15 months	Measles + mumps + rubella	Priorix®
	Chickenpox (varicella)	Varilrix®
	Haemophilus influenzae type b (Hib)	Act-HIB®
4 years	Diphtheria + tetanus + whooping cough + polio	Infanrix® IPV
11-12 years	Human papillomavirus (HPV)	Gardasil®9 (2 doses, 6 months apart)
	Tetanus + diphtheria + whooping cough	Boostrix® (from 11 years)
45 years	Tetanus + diphtheria + whooping cough	Boostrix®
65 years	Tetanus + diphtheria + whooping cough	Boostrix®
	Influenza	Brand varies
	Shingles (Zoster)	Shingrix® (2 doses, 2 - 6 months apart)

NIP8860 | Immunisation Schedule Card | 26.08.25
This resource is available from [health.govt.nz](https://www.health.govt.nz) or your local Authorised Provider

Vaccines for the National Immunisation Schedule

A reference card for vaccinators and other health professionals

From – August 2025

Pregnancy



Tdap injection (Boostrix®)
Every pregnancy, recommended from 16 weeks, funded and can be given from 13 weeks.



Influenza
1 injection (given annually)
Brand varies

6 weeks



RV1 oral vaccine (Rotarix®)



DTaP-IPV-Hep B/Hib injection (Infanrix® hexa)



PCV13 injection (Prevenar 13®)

3 months



RV1 oral vaccine (Rotarix®)



DTaP-IPV-Hep B/Hib injection (Infanrix® hexa)



MenB injection (Bexsero®)
Can be given at 8 weeks

5 months



DTaP-IPV-Hep B/Hib injection (Infanrix® hexa)



PCV13 injection (Prevenar 13®)



MenB injection (Bexsero®)
Can be given at 4 months

12 months



MMR injection (Priorix®)



PCV13 injection (Prevenar 13®)



MenB injection (Bexsero®)

15 months



MMR injection (Priorix®)



Varicella injection (Varilrix®)



Hib injection (Act-HIB®)

4 years



DTaP-IPV injection (Infanrix® IPV)

11-12 years



HPV injection (Gardasil® 9)
2 doses, 6 months apart



Tdap injection (Boostrix®)
From 11 years

45 years



Tdap injection (Boostrix®)

65 years



Tdap injection (Boostrix®)



Influenza
1 injection (given annually)
Brand varies



Shingles injection (Shingrix®)
2 doses, 2 – 6 months apart